



SUTAB[®]

(sodium sulfate, magnesium sulfate, and potassium chloride)
Tablets

1.479 g/0.225 g/0.188 g



The Dosing Regimen

SUTAB is a split-dose (2-day) regimen. A total of 24 tablets is required for complete preparation for colonoscopy. You will take the tablets in two doses of 12 tablets each. Water must be consumed with each dose of SUTAB, and additional water must be consumed after each dose.

DOSE 1—On the Day Prior to Colonoscopy

Take the tablets with water

STEP 1 Open 1 bottle of 12 tablets.

STEP 2 Fill the provided container with 16 ounces of water (up to the fill line). Swallow each tablet with a sip of water, and drink the entire amount of water over 15 to 20 minutes.



Tablets not shown actual size.

IMPORTANT: If you experience preparation-related symptoms (for example, nausea, bloating, or cramping), pause or slow the rate of drinking the additional water until your symptoms diminish.

Drink additional water

STEP 3 Approximately 1 hour after the last tablet is ingested, fill the provided container again with 16 ounces of water (up to the fill line), and drink the entire amount over 30 minutes.

STEP 4 Approximately 30 minutes after finishing the second container of water, fill the provided container with 16 ounces of water (up to the fill line), and drink the entire amount over 30 minutes.

DOSE 2—Day of the Colonoscopy


- Continue to consume only clear liquids until after the colonoscopy.
- The morning of colonoscopy (5 to 8 hours prior to the colonoscopy and no sooner than 4 hours from starting Dose 1), open the second bottle of 12 tablets.
- Repeat STEP 1 to STEP 4 from Dose 1.



Tablets not shown actual size.

IMPORTANT: You must complete all SUTAB tablets and required water at least 2 hours before colonoscopy.

Please read the full Prescribing Information and Medication Guide in the kit.

Provided as an educational service from  **Braintree**
A PART OF SEBELLA PHARMACEUTICALS

Name _____

Chart# _____

Date: _____ Dr: _____

Place: _____ T.O.A. _____

Arrange a ride to and from procedure - You will **not** be permitted to drive. Length of stay at facility - 3 to 4 hours.

7 days prior to procedure - have prescription filled at your pharmacy.

3 days before procedure - STOP the following: Vitamin E, Gingko, Ginger, Fish Oil, Garlic, Iron, Fiber Supplements, Nsaids, Ex: Ibuprofen, Motrin, Celebrex, Mobic, Naproxen/Naproxyn

2 days before procedure - Begin a low-residue diet. (See attached)

Day before procedure - Do not eat any solid food, **Clear liquid diet only.** (See attached)

* DOSE 1 - Begin SUTAB PREP @ 6:00 p.m. the night before the procedure.

"The morning of" - 6 HOURS BEFORE PROCEDURE - *DOSE 2

NOTHING TO EAT OR DRINK 4 HOURS BEFORE PROCEDURE!